

Quantum Dirac constraints, Ward identities and path integral in relativistic gauge

A.O.Barvinsky

Theory Department, Lebedev Physics Institute and Lebedev Research Center in Physics, Leninsky Prospekt 53, Moscow 117924, Russia¹

Abstract

Quantum Dirac constraints in generic constrained system are solved by directly calculating in the one-loop approximation the path integral with relativistic gauge fixing procedure. The calculations are based on the reduction algorithms for functional determinants extended to gauge theories. Explicit mechanism of transition from relativistic gauge conditions to unitary gauges, participating in the construction of this solution, is revealed by the method of Ward identities.

1. Introduction

One of the old problems in the present day high energy physics consists in the development of path integration in Dirac quantization of constrained systems – the only known regular method in problems that go beyond the scope of exactly solvable models. In contrast with non-gauge theories, the Schrodinger equation in such systems is supplemented by quantum Dirac constraints on quantum states. Moreover, in parametrized systems with a vanishing Hamiltonian there is no independent Schrodinger equation and quantum dynamics is encoded in the Dirac constraints along with their gauge invariance properties. For such problems the fundamental dynamical equations are not of a manifestly evolution type and, therefore, the path integral applications are much less straightforward.

One of the first formulations of the path integral as a solution of the non-evolutionary quantum Dirac constraints belongs to H.Leutwyler [1] who proposed a special solution of the Wheeler-DeWitt equations in the form of a naive functional integral (lacking the correct gauge-fixing procedure). The path integral in unitary gauge with exhaustive set of boundary conditions was later proposed in [2]. Then this canonical path integral was converted to the spacetime covariant form of the functional integral over Lagrangian variables and ghost fields in relativistic gauge [3, 4, 5, 6].

Of course, these results incorporate a well-known statement on equivalence of the canonical and covariant quantizations pioneered in [7]. In contrast with this, the focus

¹Electronic address: barvin@td.lpi.ac.ru

of [2, 3, 4, 5] was on nontrivial boundary conditions in spacetime. Correct treatment of these boundary conditions leads to the proof that this path integral solves quantum Dirac constraints (2.4). However, this proof [2, 3, 8, 5] has a formal nature and does not even allow one to fix the operators of these constraints which in a rather uncontrollable way depend on the calculational method for a path integral [8]. Thus, no check of the solution to quantum constraints was thus far given by direct calculations of the path integral. The goal of this paper is to perform such a check for generic systems subject to first class constraints in the one-loop approximation of semiclassical expansion.

The calculations are based on the reduction methods for one-loop functional determinants [9] extended here to the case of special boundary conditions characteristic of the coordinate representation in Dirac quantization method. These calculations lead to the Van Vleck-type solution of quantum Dirac constraints obtained earlier in the series of author's works [10, 11, 6] and as a byproduct establish the explicit mechanism of transition from relativistic gauges in the path integral to unitary gauges participating in the canonical construction of this solution.

2. Dirac quantization of constrained systems

Consider a theory with the action $S[g]$ invariant under local gauge transformations of fields $g = g^a$ with generators R_μ^a

$$R_\mu^a \frac{\delta S[g]}{\delta g^a} = 0, \quad (2.1)$$

$$S[g] = \int_{t_-}^{t_+} dt L(q, \dot{q}, N). \quad (2.2)$$

In the form (2.2), explicitly featuring time parameter, gauge fields are decomposed in two sets $g^a = (q^i(t), N^\mu(t))$ such that the Lagrangian does not involve time derivatives of $N = N^\mu(t)$ and generates the canonical action

$$S[q, p, N] = \int_{t_-}^{t_+} dt [p_i \dot{q}^i - N^\mu T_\mu(q, p)] \quad (2.3)$$

with canonical variables $(q, p) = (q^i, p_i)$ and Lagrange multipliers $N = N^\mu$ for the Dirac constraints $T_\mu(q, p)$ ². For the gauge invariant action the latter belong to the first class, i.e. satisfy the algebra $\{T_\mu, T_\nu\} = U_{\mu\nu}^\alpha T_\alpha$ of Poisson brackets with structure functions $U_{\mu\nu}^\alpha$.

² In quantum mechanical problems the range of indices $i = 1, \dots, n$ and $\mu = 1, \dots, m$ is finite and extends to infinity in field models for which we use condensed notations of two types – canonical and covariant. *Canonical* condensed indices include spatial labels and their contraction implies spatial integration. *Covariant* condensed indices involve time, and their contraction also implies time integration. They can be easily distinguished by the context in which they appear. As a rule, we imply that condensed indices are canonical when the time label is explicitly written. For Einstein theory, for example, $g^a \sim g_{\alpha\beta}(\mathbf{x}, t)$, $a = (\alpha\beta, \mathbf{x}, t)$, is the spacetime metric, $q^i = g_{ab}(\mathbf{x})$, $i = (ab, \mathbf{x})$, is a spatial metric and $N^\mu \sim g_{0\alpha}(\mathbf{x})$, $\mu = (\alpha, \mathbf{x})$ are lapse and shift functions.

Dirac quantization consists in promoting phase-space variables and constraints to the operator level $(q, p, T_\mu) \rightarrow (\hat{q}, \hat{p}, \hat{T}_\mu)$ and selecting the physical states $|\Psi\rangle$ in the representation space of $(\hat{q}, \hat{p}, \hat{T}_\mu)$ by the equation $\hat{T}_\mu |\Psi\rangle = 0$ [12, 6]. Operators $(\hat{q}, \hat{p})$ are subject to canonical commutation relations $[\hat{q}^k, \hat{p}_l] = i\hbar\delta_l^k$ and operators $\hat{T}_\mu$ should satisfy the correspondence principle with classical constraints and generalize the Poisson brackets algebra to the commutator level $[\hat{T}_\mu, \hat{T}_\nu] = i\hbar\hat{U}_{\mu\nu}^\lambda \hat{T}_\lambda$ with certain operator structure functions $\hat{U}_{\mu\nu}^\lambda$. This algebra serves as integrability conditions for quantum constraints. In the coordinate representation, $\langle q | \Psi \rangle = \Psi(q)$, $p_k = \hbar\partial/\partial q^k$, the latter become differential equations on the physical wave function $\hat{T}_\mu(q, \hbar\partial/\partial q) \Psi(q) = 0$.

Note that, without loss of generality, the nonvanishing Hamiltonian $H(q, p)$ is not included in (2.3). By extending phase space with extra canonical pair (q^0, p_0) , $q^0 \equiv t$, subject to the constraint $p_0 + H(q, p) = 0$ any system can be reduced to the case (2.3). At the quantum level this extra Dirac constraint becomes the Schrodinger equation for the wave function with parametrized time $\Psi(t, q) = \Psi(q^0, q)$. In this sense the dynamical content of any theory can be encoded in quantum constraints of the above type.

The path integral arises as a special two-point solution of these equations $\mathbf{K}(q, q')$ – the analogue of the two-point evolution operator for the Schrodinger equation [1, 2]

$$\hat{T}_\mu(q, \hbar\partial/\partial q) \mathbf{K}(q, q') = 0. \quad (2.4)$$

This is a path integral over the fields in (space)time domain $t_- < t < t_+$ with the boundary conditions related to the arguments of this kernel [1]. In view of gauge invariance this integral involves a typical Faddeev-Popov gauge fixing procedure and integration over the ghosts C and $\bar{C}$ [2, 4]

$$\mathbf{K}(q_+, q_-) = \int Dg \mu[g] DC D\bar{C} \exp \frac{i}{\hbar} \{S_{\text{gf}}[g] + \bar{C}_\mu Q_\nu^\mu C^\nu\}, \quad (2.5)$$

$$S_{\text{gf}}[g] = S[g] - \frac{1}{2} \chi^\mu c_{\mu\nu} \chi^\nu, \quad (2.6)$$

$$Q_\nu^\mu = \frac{\delta \chi^\mu}{\delta g^a} R_\nu^a, \quad (2.7)$$

$$\mu[g] = \prod_t [\det G_{ik}(t) \det c_{\mu\nu}(t)]^{1/2} \equiv (\text{Det } G_{ik} \text{Det } c_{\mu\nu})^{1/2}. \quad (2.8)$$

Here the gauge-breaking term in the gauge-fixed action $S_{\text{gf}}[g]$ is quadratic in the *relativistic* gauge conditions

$$\chi^\mu = \chi^\mu(g, \dot{g}), \quad a_\nu^\mu = -\frac{\partial \chi^\mu}{\partial \dot{N}^\nu}, \quad \det a_\nu^\mu \neq 0, \quad (2.9)$$

with some invertible gauge-fixing matrix $c_{\mu\nu}$, Q_ν^μ is a nondegenerate ghost operator and $\mu[g]$ is a local measure with $G_{ik} = \partial^2 L / \partial \dot{q}^i \partial \dot{q}^k$ – the Hessian of the classical Lagrangian³. Integration in (2.5) runs over fields with fixed values of canonical co-

³ The symbol $\det$ in (2.8) and (2.9) denotes the determinants with respect to *canonical* condensed indices in contrast with the determinants in the space of functions of time denoted by Det , defined with respect to *covariant* condensed indices.

ordinates and ghosts at $t_{\pm}$, while the boundary values of Lagrange multipliers are integrated over in the infinite range:

$$q^i(t_{\pm}) = q_{\pm}^i, \quad C^{\mu}(t_{\pm}) = 0, \quad \bar{C}_{\nu}(t_{\pm}) = 0, \quad (2.10)$$

$$-\infty < N^{\mu}(t_{\pm}) < +\infty. \quad (2.11)$$

These boundary conditions are invariant with respect to BRST transformations of the total action in (2.5) and enforce the gauge independence of the path integral [4]. They also lead to a formal proof that this integral solves quantum Dirac constraints in the coordinate representation of $q = q_+$ and $p = \hbar\partial/i\partial q_+$. This proof is based on an obvious consequence of the integration range for $N^{\mu}(t_+)$

$$\int Dg DC D\bar{C} \frac{\delta}{\delta N^{\mu}(t_+)} (\dots) = 0, \quad (2.12)$$

where ellipses denote the full integrand of the path integral (2.5). The functional differentiation here boils down to the deexponentiation of the constraint $\delta S[g]/\delta N^{\mu}(t_+) = -T_{\mu}(t_+)$ (contributions of gauge-breaking and ghost terms cancel in virtue of Ward identities [4]). This constraint can be extracted from under the integral sign in (2.12) in the form of a differential operator $\hat{T}_{\mu}(q_+, \hbar\partial/i\partial q_+)$ acting on $\mathbf{K}(q_+, q_-)$, so that the equation (2.12) takes the form of quantum Dirac constraint (2.4) [8]. This derivation is, however, purely formal and in an uncontrollable way depends on the skeletonization of the path integral [8].

3. One-loop approximation

Here we compare two semiclassical representations of $\mathbf{K}(q, q')$ – one obtained in [10, 11] by directly solving eq.(2.4) and another resulting from the one-loop approximation for the path integral (2.5). The operators $\hat{T}_{\mu}(q, \hbar\partial/i\partial q)$ closing the commutator algebra in the first subleading order of $\hbar$ -expansion were built in [11]. The solution to the corresponding quantum constraints (2.4)

$$\mathbf{K}(q, q') = \mathbf{P}(q, q') e^{\frac{i}{\hbar}\mathbf{S}(q, q')} \quad (3.1)$$

is enforced in the same approximation by the following equations for the Hamilton-Jacobi function $\mathbf{S}(q, q')$ and the preexponential factor $\mathbf{P}(q, q')$ [10, 11, 6] (see also [13] in the gravitational context)

$$T_{\mu}(q, \partial\mathbf{S}/\partial q) = 0, \quad (3.2)$$

$$\frac{\partial}{\partial q^i}(\nabla_{\mu}^i \mathbf{P}^2) = U_{\mu\lambda}^{\lambda} \mathbf{P}^2, \quad (3.3)$$

$$\nabla_{\mu}^i \equiv \left. \frac{\partial T_{\mu}}{\partial p_i} \right|_{p = \partial\mathbf{S}/\partial q}. \quad (3.4)$$

A particular solution of these equations is generated by the principal Hamilton function for $\mathbf{S}(q, q')$ – the action calculated at the classical extremal $g = g(t | q, q')$ which joins the points q and q' . This function also satisfies the Hamilton-Jacobi equations with respect to its second argument $T_\mu(q', -\partial\mathbf{S}/\partial q') = 0$. The corresponding preexponential factor is a special generalization of the Pauli-Van Vleck-Morette formula [14] in terms of the Van-Vleck matrix

$$\mathbf{S}_{ik'} = \frac{\partial^2 \mathbf{S}(q, q')}{\partial q^i \partial q^{k'}}, \quad (3.5)$$

In contrast with non-gauge theories this matrix is degenerate because it has left and right zero-eigenvalue eigenvectors [10, 11, 6]

$$\nabla_\mu^i \mathbf{S}_{ik'} = 0, \quad \mathbf{S}_{ik'} \nabla_\nu^{k'} = 0, \quad \nabla_\nu^{k'} \equiv \left. \frac{\partial T_\nu}{\partial p_{k'}} \right|_p = -\partial \mathbf{S} / \partial q'. \quad (3.6)$$

The construction of $\mathbf{P}(q, q')$ is, therefore, equivalent to the gauge-fixing procedure. It consists in adding the “gauge-breaking” term bilinear in “gauge conditions” X_i^μ and $X_{k'}^\nu$ – two sets of arbitrary covectors at the points q and q'

$$D_{ik'} = \mathbf{S}_{ik'} + X_i^\mu C_{\mu\nu} X_{k'}^\nu, \quad (3.7)$$

which replaces degenerate $\mathbf{S}_{ik'}$ by the new invertible matrix $D_{ik'}$, provided that the gauge-fixing matrix $C_{\mu\nu}$ is also invertible and these covectors produce invertible “Faddeev-Popov operators”

$$J_\nu^\mu = X_i^\mu \nabla_\nu^i, \quad J \equiv \det J_\nu^\mu, \quad (3.8)$$

$$J_\nu'^\mu = X_{i'}^\mu \nabla_\nu^{i'}, \quad J' \equiv \det J_\nu'^\mu. \quad (3.9)$$

Then the solution of the continuity equations (3.3) reads [10, 11, 6]

$$\mathbf{P}(q, q') = \left[\frac{\det D_{ik'}}{JJ' \det C_{\mu\nu}} \right]^{1/2}. \quad (3.10)$$

This solution⁴ is a direct analogue of the one-loop effective action in gauge theory [15] – the gauge field contribution $\det D_{ik'}$ partly compensated by the contribution of ghosts JJ' .

The covectors $(X_i^\mu, X_{k'}^\nu)$ are matrices of gauge conditions because, as shown in [10, 11], the quantum Hamiltonian reduction of $\mathbf{K}(q, q')$ leads to the unitary evolution operator in the physical sector defined by the *unitary* gauge conditions

$$X^\mu(q, t) = 0, \quad X_i^\mu = \frac{\partial X^\mu}{\partial q^i}. \quad (3.11)$$

Unitary gauge conditions are imposed only on phase space variables (q, p) (here only on coordinates q)⁵. In contrast with *relativistic* gauges involving Lagrange multipliers, they manifestly incorporate unitarity but lead to the loss of manifest covariance.

⁴ Particular example of such solution for a *single* Wheeler-DeWitt equation was found in [13].

⁵ Explicit time dependence of $X^\mu(q, t)$ is necessary in theories with parametrized time in order to have nontrivial time evolution with a nonvanishing physical Hamiltonian [6, 11].

On the contrary, the path integral (2.5) in relativistic gauge is potentially a space-time covariant object. Its Feynman diagrammatic technique was built in [4] with a special emphasis on boundary conditions at $t_{\pm}$, because in other respects the $\hbar$ -expansion produces a standard set of Feynman graphs. Thus, the one-loop path integral also has a form (3.1) with the same principal Hamilton function $\mathbf{S}(q, q') = S[g(t|q_+, q_-)]$ – the action at the solution $g = g(t|q_+, q_-)$ of the following boundary value problem in the gauge (2.9) [4]

$$\frac{\delta S[g]}{\delta g^a(t)} = 0, \quad \chi^\mu(g, \dot{g}) = 0, \quad t_- \leq t \leq t_+, \quad (3.12)$$

$$q(t_{\pm}) = q_{\pm}, \quad (3.13)$$

and the one-loop preexponential factor

$$\mathbf{P}(q_+, q_-) = \left(\frac{\text{Det } F_{ab}}{\text{Det } a_{ab}} \right)^{-1/2} \frac{\text{Det } Q_{\nu}^{\mu}}{\text{Det } a_{\nu}^{\mu}} \Big|_{g=g(t|q_+, q_-)}. \quad (3.14)$$

Here F_{ab} is the gauge field operator

$$F_{ab} = S_{ab} - \chi_a^{\mu} c_{\mu\nu} \chi_b^{\nu}, \quad (3.15)$$

$$S_{ab} \equiv \frac{\delta^2 S[g]}{\delta g^a \delta g^b}, \quad \chi_a^{\mu} \equiv \frac{\delta \chi^{\mu}}{\delta g^a}, \quad (3.16)$$

with the functional matrix χ_a^{μ} of linearized gauge conditions – a first-order differential operator $\chi_a^{\mu} = \chi_a^{\mu}(d/dt)\delta(t - t')$.

The determinants in (3.14) are calculated on functional spaces defined by boundary conditions for gauge and ghost operators. In relativistic gauge (2.9) these operators $F_{ab} = -a_{ab} d^2/dt^2 + \dots$ and $Q_{\nu}^{\mu} = -a_{\nu}^{\mu} d^2/dt^2 + \dots$ are of second order in time derivatives with nondegenerate matrices $a_{ab} = \partial^2 L_{\text{gf}} / \partial \dot{g}^a \partial \dot{g}^b$ (the Hessian of gauge-fixed Lagrangian) and (2.9). Their boundary conditions were derived in [4] from the conditions (2.10)-(2.11) on the integration range in the path integral. For $(n+m) \times (n+m)$ -matrix of the gauge field propagator $G^{ab} = G^{ab}(t, t')$ they form a combined set of n Dirichlet ($i = 1, \dots, n$) plus m Robin ($\mu = 1, \dots, m$) boundary conditions

$$F_{ca}(d/dt)G^{ab}(t, t') = \delta_c^b \delta(t - t'), \quad (3.17)$$

$$G^{ib}(t_{\pm}, t') = 0, \quad (3.18)$$

$$\chi_a^{\mu}(d/dt_{\pm})G^{ab}(t_{\pm}, t') = 0 \quad (3.19)$$

and Dirichlet boundary conditions for the ghost propagator $Q_{\alpha}^{-1\beta} = Q_{\alpha}^{-1\beta}(t, t')$

$$Q_{\mu}^{\alpha}(d/dt)Q_{\alpha}^{-1\beta}(t, t') = \delta_{\mu}^{\beta}, \quad Q_{\alpha}^{-1\beta}(t_{\pm}, t') = 0. \quad (3.20)$$

Conditions (3.19) belong to the Robin type because for relativistic gauges $\chi_a^{\mu}(d/dt)$ contains derivatives transversal to the boundary.

The contribution of the local measure (2.8) to $\mathbf{P}(q, q')$ in (3.14) is identically rewritten in terms of $\text{Det } a_{ab}$ and $\text{Det } a_{\nu}^{\mu}$ – functional determinants of ultralocal matrix coefficients of second derivatives in F_{ab} and Q_{ν}^{μ} . This representation will be used to show

the cancellation of strongest $\delta(0)$ -type divergences of $\text{Det } F_{ab}$ and $\text{Det } Q_\nu^\mu$ by the local measure.

Thus, the main goal of this paper consists in the proof of equality of expressions (3.10) and (3.14). This proof begins with the discussion of their gauge independence.

4. Ward identities and gauge independence

Expressions (3.10) and (3.14) are independent of the gauge choice – the matrices $(X_i^\mu, X_{k'}^\nu)$ and χ_a^μ respectively. The mechanism of this gauge independence is based on Ward identities which we consider in parallel both in canonical and covariant contexts. The Ward identity for the gauge-fixed Van Vleck matrix (3.7) follows by contracting it with ∇_μ^i and using the degeneracy relation (3.6)

$$C_{\mu\nu} X_{k'}^\nu D^{-1\ k'i} = J_\mu^{-1\ \nu} \nabla_\nu^i. \quad (4.1)$$

As a consequence, arbitrary variations of $(X_i^\mu, X_{k'}^\nu, C_{\mu\nu})$ in (3.10) vanish due to the cancellation of terms coming from $\det D_{ik'}$ and JJ' .

In the covariant (space)time context the analogue of (3.6) is the on-shell degeneracy of the functional Hessian matrix S_{ab} (3.16) – a corollary of (2.1), $R_\mu^a S_{ab} = -\delta R_\mu^a / \delta g^b \delta S / \delta g^a = 0$. It leads to the relation $R_\mu^a F_{ab} = -Q_\mu^\alpha c_{\alpha\beta} \chi_b^\beta$, which implies the Ward identity relating the gauge and ghost propagators subject to boundary conditions of the above type

$$c_{\alpha\beta} \overrightarrow{\chi}_b^\beta G^{bc}(t, t') = -Q_\alpha^{-1\ \beta}(t, t') \overleftarrow{R}_\beta^c. \quad (4.2)$$

Here the arrows show the direction in which the derivatives in differential operators $\chi_b^\beta(d/dt)$ and $R_\beta^c(d/dt')$ are acting on the arguments of Green's functions.

These Ward identities lead to gauge independence of the one-loop prefactor (3.14) provided we consistently fix the definition of functional determinants $\text{Det } F_{ab}$ and $\text{Det } Q_\nu^\mu$. Problem is that uniquely defined Green's functions do not yet uniquely fix these determinants. Variational equations for the latter involve the functional composition of Green's functions with variations of their operators: $G^{ba} \delta F_{ab}$ and $Q_\mu^{-1\ \nu} \delta Q_\nu^\mu$. However, kernels of propagators are not smooth functions and their irregularity enhances when they are acted upon by the derivatives of $\delta F_{ab}(d/dt)$ and $\delta Q_\nu^\mu(d/dt)$. Therefore, one has to prescribe the way how these derivatives act on both arguments of Green's functions. We assume that the varied differential operators $\delta F_{ab}(d/dt)$ and $\delta Q_\nu^\mu(d/dt)$ are understood as acting in two different ways

$$\delta \ln \text{Det } F_{ab} = G^{ba} \overleftrightarrow{\delta} F_{ab}, \quad (4.3)$$

$$\delta \ln \text{Det } Q_\nu^\mu = Q_\mu^{-1\ \nu} \overleftarrow{\delta} Q_\nu^\mu. \quad (4.4)$$

In contrast with the ghost operator, for which both of its derivatives are acting on *one* argument of the Green's function, eq.(4.3) here implies a symmetric action of $\delta F_{ab}(d/dt)$

on *both* arguments of $G^{ba}(t, t')$ in the sense that $L_{\text{gf}}^{(2)} = (1/2)\varphi^a(t) \overleftrightarrow{F}_{ab}(d/dt) \varphi^b(t)$ represents the quadratic part of the gauge-fixed Lagrangian in perturbations of field variables φ^a [9]. It contains the squares of $\dot{\varphi}^a(t)$ rather than $\ddot{\varphi}^a(t)$ (and $\delta\overleftrightarrow{F}_{ab}(d/dt)$ obviously implies arbitrary variation of this kernel).

With these conventions the gauge variation of gauge and ghost determinants in (3.14)

$$\delta_\chi \ln \text{Det } F_{ab} = -2 c_{\alpha\beta} \overrightarrow{\chi}_b^\beta G^{ba} \delta\overleftarrow{\chi}_a^\alpha, \quad (4.5)$$

$$\delta_\chi \ln \text{Det } Q_\nu^\mu = Q_\alpha^{-1\beta} \overleftarrow{R}_\beta^b \delta\overleftarrow{\chi}_b^\alpha \quad (4.6)$$

cancel out in virtue of (4.2). This proves the gauge independence of (3.14) along with fixing the prescription for the variational definition (4.3)-(4.4) of the functional determinants⁶.

5. Reduction algorithms for functional determinants

Equality of expressions (3.10) and (3.14) follows from the reduction algorithms for functional determinants which reduce their dimensionality from the functional dimensionality of Det 's in (3.14) to that of det 's in (3.10) [9]. The simplest example of such algorithms is the relation (2.8) for a purely divergent local measure. For an ultralocal operator $a_{ab} = a_{ab}(t) \delta(t - t')$ one has

$$\ln \text{Det } a_{ab} = \delta(0) \int_{t_-}^{t_+} dt \ln \text{det } a_{ab}(t). \quad (5.1)$$

For differential operators the reduction algorithms result from the functional integration of eqs.(4.3)-(4.4). For symmetric operators with Dirichlet boundary conditions they were considered in much detail in [9]. Here we generalize them to the case of combined Dirichlet-Robin boundary conditions (3.18)-(3.19) for F_{ab} and to the case of non-symmetric ghost operator Q_ν^μ .

Starting with $F_{ab}(d/dt)$, we note that this operator satisfies the Wronskian relation with arbitrary test functions $\varphi_{1,2}^a$

$$\varphi^a (F_{ab} \varphi^b) - (F_{ba} \varphi^a) \varphi^b = -\frac{d}{dt} \left[\varphi^a (W_{ab} \varphi^b) - (W_{ba} \varphi^a) \varphi^b \right], \quad (5.2)$$

where $W_{ab} = a_{ab} d/dt + \dots$ is the *Wronskian* operator. It participates in the variational equation for the canonical momentum, $\delta(\partial L_{\text{gf}}/\partial \dot{g}) = W(d/dt) \delta g(t)$, and, in particular, has a special form of its μ -component given by the operator of linearized gauge conditions $W_{\mu b}(d/dt) = a_\mu^\alpha c_{\alpha\beta} \chi_b^\beta(d/dt)$.

⁶ In context of the one-loop effective action Ward identities were considered in [15]. Here the main emphasis in their mechanism is focused on boundary conditions and accurate definition of the functional determinants. Another choice of these variational equations leads to extra surface terms violating gauge independence.

Similarly to [9], introduce for $F_{ab}(d/dt)$ two complete sets of basis functions $u_-^a(t)$ and $u_+^a(t)$

$$F_{ab}(d/dt) u_{\pm}^b(t) = 0, \quad (5.3)$$

$$u_+^i(t_+) = 0, \quad \vec{W}_{\mu a} u_+^a(t_+) = 0, \quad (5.4)$$

$$u_-^i(t_-) = 0, \quad \vec{W}_{\mu a} u_-^a(t_-) = 0. \quad (5.5)$$

satisfying the boundary conditions (3.18)-(3.19) respectively at t_- and t_+ , the Robin conditions (3.19) being rewritten in terms of $W_{\mu b}(d/dt)$. We assume that the indices enumerating complete sets of these basis functions are encoded in subscripts $\pm$. Then, the t -independent Wronskian inner product of basis functions forms a matrix in vector space of these indices. In view of boundary conditions it has only two nonvanishing blocks given by two mutually transposed matrices

$$\Delta_{-+} = u_-^a (W u_+)_a - (W u_-)_a u_+^b, \quad \Delta_{+-} = -\Delta_{-+}^T, \quad (5.6)$$

where for brevity we introduced the notation $(W u_{\pm})_a \equiv \vec{W}_{ab} u_{\pm}^b$. In terms of basis functions the Green's function of the mixed Dirichlet-Robin boundary value problem (3.17)-(3.19) has the following representation

$$G^{ab}(t, t') = -\theta(t-t') u_+^a(t) (\Delta_{-+})^{-1} u_-^b(t') + \theta(t'-t) u_-^a(t) (\Delta_{+-})^{-1} u_+^b(t'), \quad (5.7)$$

where $\theta(x)$ is the step function: $\theta(x) = 1$ for $x > 0$ and $\theta(x) = 0$ for $x < 0$.

After substituting (5.7) to (4.3) the $\delta(0)$ -type terms, which originate from the derivatives of θ -functions, reduce to the variation of (5.1) [9], so that the strongest divergences of $\text{Det } F_{ab}$ are cancelled by the local measure [16]. The rest can be transformed by integrating by parts and using the equations (5.3) or their corollary $F_{ab} u_{\pm}^b = -F_{ab} \delta u_{\pm}^b$. This allows one in a systematic way to reduce the full answer to surface terms at $t = t_{\pm}$ which form the total variation that can easily be functionally integrated to give (see [17] for details)

$$\left(\frac{\text{Det } \overset{\leftrightarrow}{F}_{ab}}{\text{Det } a_{ab}} \right)^{-1/2} = \text{const} \left(\frac{\det (\mathbf{S}_{ik'} + X_i^{\mu} C_{\mu\nu} X_{k'}^{\nu})}{\det C_{\mu\nu}} \right)^{1/2}, \quad (5.8)$$

where the matrices $(X_i^{\mu}, X_{k'}^{\nu}, C_{\mu\nu})$ in terms of basis functions of the above type read

$$X_i^{\nu} = (W u_+)_i(t_+) \Delta_{-+}^{-1} u_-^{\nu}(t_-), \quad (5.9)$$

$$X_{k'}^{\mu} = u_+^{\mu}(t_+) \Delta_{-+}^{-1} (W u_-)_k(t_-), \quad (5.10)$$

$$C^{\mu\nu} = u_+^{\mu}(t_+) \Delta_{-+}^{-1} u_-^{\nu}(t_-), \quad C_{\mu\nu} = (C^{\mu\nu})^{-1} \quad (5.11)$$

and $\mathbf{S}_{ik'}$ arising here as $\mathbf{S}_{ik'} = -(W u_+)_i(t_+) \Delta_{-+}^{-1} (W u_-)_k(t_-)$ coincides with the Van Vleck matrix (3.5) for the principal Hamilton function [17]. This equation is a generalization of the results of paper [9] extending the path-integral derivation of the Pauli-Van

Vleck-Morette formula [14] to gauge theories. Its comparison with (3.7),(3.10) shows that the quantities (5.9)-(5.11) are special unitary gauge conditions and gauge-fixing matrix induced by the relativistic gauge fixing procedure.

The functional integration of eq.(4.4) for $\text{Det } \vec{Q}_\nu^\mu$ repeats the calculations of [9] with modifications caused by the asymmetry of the ghost operator. This asymmetry results in a double set of right and left basis functions which give rise to the representation of $Q_\alpha^{-1\beta}$ similar to (5.7) and after functional integration yield the analogue of the Pauli-Van Vleck formula with a special Van Vleck matrix [17]. In view of Ward identity (4.2) special combinations of *ghost* field basis functions entering this matrix can be related to the unitary gauge conditions (5.9)-(5.10) built in terms of the basis functions of the *gauge* operator. The final reduction algorithm then reads (see [17] for details)

$$\frac{\text{Det } \vec{Q}_\nu^\mu}{\text{Det } a_\nu^\mu} = \text{const} (\det J_\nu^\mu \det J_\nu'^\mu)^{-1/2}, \quad (5.12)$$

where the matrices J_ν^μ and $J_\nu'^\mu$ coincide with unitary Faddeev-Popov operators (3.8)-(3.9) in unitary gauges of the above type.

The substitution of (5.8) and (5.12) to (3.14) accomplishes the proof of the needed equality of (3.10) and (3.14).

6. Conclusions

The virtue of equivalence of (3.10) and (3.14) is that it establishes the explicit mechanism of transition from relativistic to unitary gauge conditions. The relations between them are nonlocal in time – the matrices of unitary gauge-fixing procedure (5.9)-(5.11) express in terms of basis functions of the gauge field operator, nonlocally depending on its relativistic gauge. Such a transition is very important because it proves intrinsic unitarity of a manifestly covariant quantization in terms of a Lagrangian path integral, and the mechanism of this transition revealed here is free from singularities inherent to the usual ϵ -procedure of formally identical transformations in the path integral [7].

In physical applications, Feynman diagrammatic expansion of the path integral as a means of solving *noncovariant* quantum Dirac constraints [19] is of crucial importance due to *spacetime covariance* of their solution that can be attained by a suitable choice of relativistic gauge conditions. Important implications of this technique belong to quantum cosmology of the early universe [18] where correct predictions can be achieved only within spacetime covariant approach to loop effects.

The aspects of gauge independence considered above are important for problems in spacetimes with boundaries or nontrivial time foliations. There exists a long list of gauge-dependent results for a formally gauge independent quantity – the one-loop effective action [20]. Exhaustive explanation of these discrepancies can be expected on the basis of Ward identities with a special emphasis on boundary conditions.

Finally, the physics of wormholes in Euclidean quantum gravity [21] also belongs to the scope of our result. The predictions of this theory are based, in particular,

on the existence of a negative mode on the wormhole instanton [22] – a formal extrapolation of the mechanism applicable only to non-gravitational systems [23]. Thus, these predictions should be revised from the viewpoint of the Wheeler-DeWitt equation [24, 22] – the gravitational quantum Dirac constraint. This negative mode belongs to the *nondynamical* conformal sector, and in the Lorentzian theory its contribution is cancelled by ghost fields in relativistic gauges. Therefore, one should expect a similar cancellation by the mechanism of Ward identities also in Euclidean theory. Other problems related to the indefiniteness of the Euclidean gravitational action include the lack of strong ellipticity of the Dirichlet-Robin boundary value problem (3.17)-(3.19) [25]. The proposed technique is a direct avenue towards the resolution of these issues which are currently under study.

Acknowledgements

This work was supported by the Russian Foundation for Basic Research under grants 96-02-16287 and 96-02-16295 and the European Community Grant INTAS-93-493-ext. Partly this work has been made possible also due to the support by the Russian Research Project “Cosmomicrophysics”.

References

- [1] H.Leutwyler, Phys. Rev. **134** (1964) B1155
- [2] A.O.Barvinsky, in Proceedings of the 9th Conference on General Relativity and Gravitation (Jena, 1980) p. 540; A.O.Barvinsky and V.N.Ponomariov, Phys.Lett. **B167** (1986) 289
- [3] J.B.Hartle and S.W.Hawking, Phys.Rev. **D28** (1983) 2960
- [4] A.O.Barvinsky, Phys.Lett. **B195** (1987) 344
- [5] J.J.Halliwell and J.B.Hartle, Phys.Rev. **D43** (1991) 1170
- [6] A.O.Barvinsky, Phys. Rep. **230** (1993) 237
- [7] E.S.Fradkin and G.A.Vilkovisky, Phys.Lett. **B55** (1975) 224, CERN Report TH-2332 (1977)
- [8] A.O.Barvinsky, Phys.Lett. **B175** (1987) 401
- [9] A.O.Barvinsky, Phys. Rev. **D50** (1994) 5115
- [10] A.O.Barvinsky, Phys.Lett. **B241** (1990) 201
- [11] A.O.Barvinsky and V.Krykhtin, Class.Quantum Grav. **10** (1993) 1957

- [12] M.Henneaux, Phys. Rep. **126** (1985) 1; I.A.Batalin and E.S.Fradkin, Riv.Nuovo Cim. **9** (1986) 1, Ann. Inst. Henri Poincare **49** (1988) 145; I.A.Batalin and R.Marnelius, Nucl.Phys. **B442** (1995) 669
- [13] C.Kiefer and T.P.Singh, Phys.Rev. **D44** (1991) 106; C.Kiefer, preprint gr-qc/9312015
- [14] C.DeWitt-Morette, Ann.Phys. **97** (1976) 367
- [15] A.O.Barvinsky and G.A.Vilkovisky, Phys.Rep. **119** (1985) 1
- [16] E.S.Fradkin and G.A.Vilkovisky, On Renormalization of Quantum Field Theory in Curved Spacetime, Institute for Theoretical Physics, University of Berne preprint (1976)
- [17] A.O.Barvinsky, Solution of quantum Dirac constraints via path integral, preprint FIAN/TD/97-21, hep-th/9711164
- [18] A.O.Barvinsky, A.Yu.Kamenshchik and I.V.Mishakov, Nucl.Phys. **B491** (1997) 387
- [19] A.O.Barvinsky and C.Kiefer, Wheeler-DeWitt equation and Feynman diagrams, preprint FIAN/TD/97-20, gr-qc/9711037 (1997)
- [20] P.A. Griffin and D.A. Kosower, Phys.Lett. **B233** (1989) 295; G.Esposito, A.Yu.Kamenshchik, I.V.Mishakov and G.Pollifrone, Phys.Rev. **D52** (1995) 3457
- [21] G.V.Lavrelashvili, V.A.Rubakov and P.G.Tinyakov, JETP Lett. **46** (1987) 167; S.B.Giddings and A.Strominger, Nucl. Phys. **B306** (1988) 890; S.Coleman, Nucl. Phys. **B307** (1988) 867
- [22] V.A.Rubakov and O.Yu.Shvedov, Phys.Lett. **B383** (1996) 258
- [23] S.Coleman, Nucl.Phys. **B298** (1988) 178
- [24] T.Tanaka and M.Sasaki, Prog.Theor.Phys. **88** (1992) 503
- [25] I.G.Avramidy and G.Esposito, Lack of strong ellipticity in Euclidean quantum gravity, preprint DSF 97/43, hep-th/9708163